

Copyright © 2016 Moradigaravand et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

1 **New data letter**2 ***DfrA-thyA* double deletion in *para*-aminosalicylic acid resistant *Mycobacterium tuberculosis* Beijing**  
3 **strains**

4

5 Running title: PAS resistance due to *dfrA-thyA* deletion

6

7 Danesh Moradigaravand<sup>1#</sup>, Louis Grandjean<sup>2-4#</sup>, Elena Martinez<sup>5-7#</sup>, Hao Li<sup>8#</sup>, Jun Zheng<sup>9#</sup>, Jorge  
8 Coronel<sup>4</sup>, David Moore<sup>3,4</sup>, M. Estée Török<sup>10-12</sup>, Vitali Sintchenko<sup>5-7</sup>, Hairong Huang<sup>13</sup>, Babak Javid<sup>8,10</sup>,  
9 Julian Parkhill<sup>1</sup>, Sharon J. Peacock<sup>1,3,10</sup> & Claudio U. Köser<sup>10\*</sup>

10

11 <sup>1</sup>Wellcome Trust Sanger Institute, Hinxton, UK12 <sup>2</sup>Wellcome Centre for Clinical Tropical Medicine, Imperial College London, St Mary's Campus, London,  
13 UK14 <sup>3</sup>London School of Hygiene and Tropical Medicine, London, UK15 <sup>4</sup>Laboratorio de Investigación en Enfermedades Infecciosas, Universidad Peruana Cayetano Heredia,  
16 Lima, Peru17 <sup>5</sup>Centenary Institute and Marie Bashir Institute for Infectious Diseases and Biosecurity, The University  
18 of Sydney, Sydney, Australia19 <sup>6</sup>NSW Mycobacterium Reference Laboratory, Centre for Infectious Diseases and Microbiology  
20 Laboratory Services, Institute of Clinical Pathology and Medical Research—Pathology West, Sydney,  
21 Australia22 <sup>7</sup>Centre for Infectious Diseases and Microbiology—Public Health, Westmead Hospital, Western  
23 Sydney Local Health District, Sydney, Australia24 <sup>8</sup>Collaborative Innovation Centre for the Diagnosis and Treatment of Infectious Diseases, School of  
25 Medicine, Tsinghua University, Beijing, China26 <sup>9</sup>Faculty of Health Sciences, University of Macau, Macau SAR, China27 <sup>10</sup>Department of Medicine, University of Cambridge, Cambridge, UK28 <sup>11</sup>Department of Infectious Diseases, Cambridge University Hospitals NHS Foundation Trust,  
29 Cambridge, UK

30 <sup>12</sup>Clinical Microbiology and Public Health Laboratory, Public Health England, Cambridge, UK

31 <sup>13</sup>National Clinical Laboratory on Tuberculosis, Beijing Key laboratory on Drug-resistant Tuberculosis

32 Research, Beijing Chest Hospital, Capital Medical University, Beijing Tuberculosis and Thoracic Tumor

33 Institute, Beijing, China

34

35 <sup>#</sup>Contributed equally

36 \*Corresponding author. Tel: +44-1223-331664; Fax: +44-1223-336846; E-mail: cuk21@cam.ac.uk

37

38 Keywords: *Mycobacterium tuberculosis*, *para*-aminosalicylic acid resistance, dihydrofolate reductase,

39 thymidylate synthase

40

41 *Para*-aminosalicylic acid (PAS) is a group 4 anti-tuberculosis agent (1). It targets folate metabolism as

42 shown in Fig. S1, which also summarises the known resistance mechanisms to this pro-drug (2).

43 Recently, we reported a multidrug-resistant (MDR) *Mycobacterium tuberculosis* Beijing strain

44 harbouring a deletion of both *dfrA* and *thyA* from Australia (Fig. 1A and Table S1) (3). Since then, we

45 have found deletions affecting both genes in five further MDR Beijing strains (two isolated in Australia

46 and three from Peru) and one extensively drug-resistant (XDR) Beijing strain from China. The

47 Australian MDR strains were recovered from three patients with no apparent epidemiological links

48 and were likely infected in their country of origin (Table S1). The three Peruvian isolates were closely

49 related and consequently shared the same deletion, whereas the remaining strains were distantly

50 related and had deletions that differed in size (Fig. 1A). Consequently, these five distinct deletions

51 were acquired independently, which can be a signal for positive selection of resistance mechanisms.

52 In line with this hypothesis, the strains from Australia and China were PAS resistant when tested with

53 the BACTEC MGIT 960 system and on Löwenstein-Jensen medium, respectively (Supplementary

54 Methods). Two out of the three Peruvian deletion mutants were also PAS resistant on 7H10 medium

55 at 8 µg/mL, whereas the two closely related ancestral wild-type strains were susceptible (Fig. 1B). We

56 were unable to retest the strains at 2 µg/mL, the recommended critical concentration by the Clinical

57 and Laboratory Standards Institute and World Health Organization, which would have clarified

58 whether the susceptible result for the third deletion mutant was an artefact (1, 4).

59           The observation that *dfrA* could be deleted was remarkable in light of our current  
60 understanding of folate metabolism in *M. tuberculosis*. Two studies suggested that *dfrA* is essential *in*  
61 *vitro* in the H37Rv laboratory strain (5, 6). More recently, it was shown that *dfrA* is conditionally  
62 essential and can only be knocked out in H37Rv if Rv2671 is over-expressed *in trans*, presumably due  
63 to its much lower dihydrofolate reductase activity compared with DfrA (7, 8). Our *in silico* analysis of  
64 the seven *dfrA-thyA* double deletion mutants did not reveal any known Rv2671 mutations (Table S1),  
65 such as the G to A upstream mutation at position -12 that results in its over-expression and  
66 consequently confers PAS resistance (this mutation was incorrectly referred to as affecting position -  
67 11 in two of our prior studies (7, 9)). Assuming that no other pertinent differences exist that are  
68 specific to the Beijing genotype relative to H37Rv or that a yet unknown acquired mutation elsewhere  
69 in the genome were present that resulted in the over-expression of Rv2671, we propose that this  
70 apparent contradiction can be reconciled if the essentiality of *dfrA* was not only dependent on the  
71 expression level of Rv2671, but also on the presence of wild-type *thyA*. The fact that *thyA* was deleted  
72 in all seven *dfrA* mutants meant that only the second thymidylate synthase, encoded by the essential  
73 *thyX*, was active in these strains (Fig. S1). Contrary to ThyA, ThyX generates tetrahydrofolate rather  
74 than dihydrofolate upon catalysis and therefore does not require high dihydrofolate reductase  
75 activity to provide sufficient levels of tetrahydrofolate (2). This is in line with the fact that *dfrA* is not  
76 required in bacterial species that lack *thyA* (10). Consequently, Rv2671 appeared to be sufficient to  
77 sustain growth, even without being over-expressed in these deletion mutants. It should therefore be  
78 possible to knock out *dfrA* in strains of *M. tuberculosis* with inactive *thyA*. Moreover, the adjacent  
79 location of *thyA* and *dfrA* in genome should make their simultaneous deletion possible (Fig. 1A).

80           Interestingly, all but one of the deletion mutants also convergently acquired mutations  
81 upstream of *thyX* compared to the two closely related Peruvian control strains (Fig. 1B & Table S1)  
82 (11). In fact, the cluster of three Peruvian strains and two of the unrelated Australian strains shared  
83 the same C to T upstream mutation at position -16 that has been previously found to be associated  
84 with resistance to several drugs and experimentally shown to result in the over-expression of *thyX*  
85 (12). It is therefore plausible that these changes compensated for the reduced expression levels and  
86 enzymatic activity of ThyX compared to ThyA (11, 13). Based on our data, however, it was not

87 possible to deduce whether the *thyX* mutations were acquired after the deletions of *thyA* and *dfrA* in  
88 each strain, as would be expected with compensatory mutations (11).

89 In summary, these data demonstrated that the folate metabolism and the genetic basis of  
90 PAS resistance are more complex than previously appreciated, which is relevant for the development  
91 of novel DfrA and ThyX inhibitors and potentially the use of trimethoprim-sulfamethoxazole to treat  
92 drug-resistant tuberculosis (Fig. S1) (14-25). Although deletions are often excluded from large-scale  
93 whole genome studies owing to the limited read lengths next-generation sequencers and the fact that  
94 algorithms are optimised for SNP calling, this study highlighted that deletions can no longer be  
95 ignored (3, 26).

96

97 **Fig. 1.** Analysis of *dfrA* and *thyA* deletion strains, all of which tested PAS resistant, with the exception  
98 of PH107\_CA033M\_1. (a) Diagram of deletions in seven clinical strains compared with the wild-type  
99 H37Rv laboratory strain. The scale at the top corresponds to the genome position in H37Rv. The letter  
100 in parentheses denotes the country of isolation (Australia (A), China (C), and Peru (P)). Mtb97 was  
101 reported previously (3). (b) Maximum likelihood tree based on whole-genome data of the three  
102 Peruvian deletion mutants, which also share a mutation upstream of *thyX* that is also present in  
103 Mtb97 and Mtb78 (Table S1), and two closely related wild-type strains, which were PAS susceptible.

104     **Funding information**

105     This publication presents independent research supported by the Health Innovation Challenge Fund  
106     (HICF-T5-342 and WT098600), a parallel funding partnership between the UK Department of Health  
107     and Wellcome Trust, and grant SRG2015-00006-FHS from the University of Macau. The views  
108     expressed in this publication are those of the authors and not necessarily those of the Department of  
109     Health, Public Health England, or the Wellcome Trust. M. E. T. is a Clinician Scientist Fellow funded by  
110     the Academy of Medical Sciences and the Health Foundation and the NIHR Cambridge Biomedical  
111     Research Centre. E.M. was supported by the Australian National Health and Medical Research  
112     Council's Centre for Research Excellence in Tuberculosis. C. U. K. is a Junior Research Fellow at  
113     Wolfson College, Cambridge.

114

115     **Acknowledgements**

116     We would like to thank L. Camacho for helpful comments regarding this topic.

117

118     **Conflicts of interest**

119     M. E. T., J. P., S. J. P. and C. U. K. have collaborated with Illumina Inc. on a number of scientific  
120     projects. M. E. T. and S. J. P. have received funding for travel and accommodation from Illumina Inc. J.  
121     P. has received funding for travel and accommodation from Pacific Biosciences Inc. and Illumina Inc.  
122     B. J. is a Tsinghua-Janssen Scholar. C. U. K. is a consultant for the Foundation for Innovative New  
123     Diagnostics and was a technical advisor for the Tuberculosis Guideline Development Group of the  
124     World Health Organization. The Bill & Melinda Gates Foundation and Janssen Pharmaceutica covered  
125     C. U. K.'s travel and accommodation to present at meetings. The European Society of  
126     Mycobacteriology awarded C. U. K. the Gertrud Meissner Award, which is sponsored by Hain  
127     Lifescience. All other authors: none to declare.

128

129

130     **References**

- 131     1.     **World Health Organization.** Companion handbook to the WHO guidelines for the  
132     programmatic management of drug-resistant tuberculosis (2014).

- 133 [http://apps.who.int/iris/bitstream/10665/130918/1/9789241548809\\_eng.pdf?ua=1&ua=1](http://apps.who.int/iris/bitstream/10665/130918/1/9789241548809_eng.pdf?ua=1&ua=1)  
134 (accessed 13.8.2015).
- 135 2. **Minato Y, Thiede JM, Kordus SL, McKlveen EJ, Turman BJ, Baughn AD.** 2015.  
136 *Mycobacterium tuberculosis* folate metabolism and the mechanistic basis for *para*-  
137 aminosalicylic acid susceptibility and resistance. *Antimicrob Agents Chemother* **59**:5097-  
138 5106.
- 139 3. **Martinez E, Holmes N, Jelfs P, Sintchenko V.** 2015. Genome sequencing reveals novel  
140 deletions associated with secondary resistance to pyrazinamide in MDR *Mycobacterium*  
141 *tuberculosis*. *J Antimicrob Chemother* **70**:2511-2514.
- 142 4. **Clinical and Laboratory Standards Institute.** 2011. Susceptibility testing of mycobacteria,  
143 nocardiae, and other aerobic actinomycetes, 2nd edition Approved standard. CLSI document  
144 M24-A2.
- 145 5. **Griffin JE, Gawronski JD, Dejesus MA, Ioerger TR, Akerley BJ, Sassetti CM.** 2011. High-  
146 resolution phenotypic profiling defines genes essential for mycobacterial growth and  
147 cholesterol catabolism. *PLoS Pathog* **7**:e1002251.
- 148 6. **Sassetti CM, Boyd DH, Rubin EJ.** 2003. Genes required for mycobacterial growth defined by  
149 high density mutagenesis. *Mol Microbiol* **48**:77-84.
- 150 7. **Zheng J, Rubin EJ, Bifani P, Mathys V, Lim V, Au M, Jang J, Nam J, Dick T, Walker JR, Pethe**  
151 **K, Camacho LR.** 2013. *Para*-aminosalicylic acid is a prodrug targeting dihydrofolate reductase  
152 in *Mycobacterium tuberculosis*. *J Biol Chem* **288**:23447-23456.
- 153 8. **Cheng YS, Sacchettini JC.** 2016. Structural insights into *Mycobacterium tuberculosis* Rv2671  
154 protein as a dihydrofolate reductase functional analogue contributing to *para*-aminosalicylic  
155 acid resistance. *Biochemistry* **55**:1107-1119.
- 156 9. **Zhang X, Liu L, Zhang Y, Dai G, Huang H, Jin Q.** 2015. Genetic determinants involved in *p*-  
157 aminosalicylic acid resistance in clinical isolates from tuberculosis patients in northern China  
158 from 2006 to 2012. *Antimicrob Agents Chemother* **59**:1320-1324.
- 159 10. **Mylykallio H, Leduc D, Filee J, Liebl U.** 2003. Life without dihydrofolate reductase Foa.  
160 *Trends Microbiol* **11**:220-223.

- 161 11. Merker M, Kohl TA, Roetzer A, Truebe L, Richter E, Rüsche-Gerdes S, Fattorini L, Oggioni MR,  
162 Cox H, Varaine F, Niemann S. 2013. Whole genome sequencing reveals complex evolution  
163 patterns of multidrug-resistant *Mycobacterium tuberculosis* Beijing strains in patients. PLoS  
164 One 8:e82551.
- 165 12. Zhang H, Li D, Zhao L, Fleming J, Lin N, Wang T, Liu Z, Li C, Galwey N, Deng J, Zhou Y, Zhu Y,  
166 Gao Y, Wang T, Wang S, Huang Y, Wang M, Zhong Q, Zhou L, Chen T, Zhou J, Yang R, Zhu G,  
167 Hang H, Zhang J, Li F, Wan K, Wang J, Zhang XE, Bi L. 2013. Genome sequencing of 161  
168 *Mycobacterium tuberculosis* isolates from China identifies genes and intergenic regions  
169 associated with drug resistance. Nat Genet 45:1255-1260.
- 170 13. Fivian-Hughes AS, Houghton J, Davis EO. 2012. *Mycobacterium tuberculosis* thymidylate  
171 synthase gene *thyX* is essential and potentially bifunctional, while *thyA* deletion confers  
172 resistance to *p*-aminosalicylic acid. Microbiology 158:308-318.
- 173 14. Kögler M, Busson R, De Jonghe S, Rozenski J, Van Belle K, Louat T, Munier-Lehmann H,  
174 Herdewijn P. 2012. Synthesis and evaluation of 6-aza-2'-deoxyuridine monophosphate  
175 analogs as inhibitors of thymidylate synthases, and as substrates or inhibitors of thymidine  
176 monophosphate kinase in *Mycobacterium tuberculosis*. Chem Biodivers 9:536-556.
- 177 15. Kumar A, Zhang M, Zhu L, Liao RP, Mutai C, Hafsati S, Sherman DR, Wang MW. 2012. High-  
178 throughput screening and sensitized bacteria identify an *M. tuberculosis* dihydrofolate  
179 reductase inhibitor with whole cell activity. PLoS One 7:e39961.
- 180 16. Vilchèze C, Jacobs WR, Jr. 2012. The combination of sulfamethoxazole, trimethoprim, and  
181 isoniazid or rifampin is bactericidal and prevents the emergence of drug resistance in  
182 *Mycobacterium tuberculosis*. Antimicrob Agents Chemother 56:5142-5148.
- 183 17. Parchina A, Froeyen M, Margamuljana L, Rozenski J, De Jonghe S, Briers Y, Lavigne R,  
184 Herdewijn P, Lescrinier E. 2013. Discovery of an acyclic nucleoside phosphonate that inhibits  
185 *Mycobacterium tuberculosis* ThyX based on the binding mode of a 5-alkynyl substrate  
186 analogue. ChemMedChem 8:1373-1383.
- 187 18. McGuigan C, Derudas M, Gonczy B, Hinsinger K, Kandil S, Pertusati F, Serpi M, Snoeck R,  
188 Andrei G, Balzarini J, McHugh TD, Maitra A, Akorli E, Evangelopoulos D, Bhakta S. 2014.



- 189 ProTides of N-(3-(5-(2'-deoxyuridine))prop-2-ynyl)octanamide as potential anti-tubercular  
190 and anti-viral agents. *Bioorg Med Chem* **22**:2816-2824.
- 191 19. **Alexandrova LA, Chekhov VO, Shmalenyuk ER, Kochetkov SN, El-Asrar RA, Herdewijn P.**  
192 2015. Synthesis and evaluation of C-5 modified 2'-deoxyuridine monophosphates as  
193 inhibitors of *M. tuberculosis* thymidylate synthase. *Bioorg Med Chem* **23**:7131-7137.
- 194 20. **Hong W, Wang Y, Chang Z, Yang Y, Pu J, Sun T, Kaur S, Sacchettini JC, Jung H, Lin Wong W,**  
195 **Fah Yap L, Fong Ngeow Y, Paterson IC, Wang H.** 2015. The identification of novel  
196 *Mycobacterium tuberculosis* DHFR inhibitors and the investigation of their binding  
197 preferences by using molecular modelling. *Sci Rep* **5**:15328.
- 198 21. **Lele AC, Raju A, Khambete MP, Ray MK, Rajan MG, Arkile MA, Jadhav NJ, Sarkar D, Degani**  
199 **MS.** 2015. Design and synthesis of a focused library of diamino triazines as potential DHFR  
200 inhibitors. *ACS Med Chem Lett* **6**:1140-1144.
- 201 22. **Mugumbate G, Abrahams KA, Cox JA, Papadatos G, van Westen G, Lelievre J, Calus ST,**  
202 **Loman NJ, Ballell L, Barros D, Overington JP, Besra GS.** 2015. Mycobacterial dihydrofolate  
203 reductase inhibitors identified using chemogenomic methods and *in vitro* validation. *PLoS*  
204 *One* **10**:e0121492.
- 205 23. **Raju A, Degani MS, Khambete MP, Ray MK, Rajan MG.** 2015. Antifolate activity of plant  
206 polyphenols against *Mycobacterium tuberculosis*. *Phytother Res* **29**:1646-1651.
- 207 24. **Singh V, Brecik M, Mukherjee R, Evans JC, Svetlikova Z, Blasko J, Surade S, Blackburn J,**  
208 **Warner DF, Mikusova K, Mizrahi V.** 2015. The complex mechanism of antimycobacterial  
209 action of 5-fluorouracil. *Chem Biol* **22**:63-75.
- 210 25. **Tawari NR, Bag S, Raju A, Lele AC, Bairwa R, Ray MK, Rajan MG, Nawale LU, Sarkar D,**  
211 **Degani MS.** 2015. Rational drug design, synthesis and biological evaluation of dihydrofolate  
212 reductase inhibitors as antituberculosis agents. *Future Med Chem* **7**:979-988.
- 213 26. **Machado D, Couto I, Perdigao J, Rodrigues L, Portugal I, Baptista P, Veigas B, Amaral L,**  
214 **Viveiros M.** 2012. Contribution of efflux to the emergence of isoniazid and multidrug  
215 resistance in *Mycobacterium tuberculosis*. *PLoS One* **7**:e34538.
- 216



